

PAPILIO AMPHITRYON.



CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatæ.

Alæ (sedentis) erectæ sursumque conniventæ
(volatu diurno.)

CHARACTER SPECIFICUS, &c.

PAPILIO alis anticis apice flavo-maculatis, posticis fascia flavescente, subtus lunulis cæruleis.

PAPILIO AMPHITRION. P. alis nigris fascia inæquali flava, posterioribus subtus striga e punctis flavis lunulisque cæruleis.

Lin. Syst. Nat. Gmel. p. 2247.

PAPILIO AMPHITRYON.

Cram. t. 7. f. A. B.

Eq. Archiv.

Americæ Australis partes calidiores incolit Papilio Amphitryon, coloribus interdum leviter varians. Magnitudinem veram ostendit tabula.

600
49



Pyr

AMPHITRYON.

GENERIC CHARACTER, &c.

Antennæ thickening towards the upper part, and generally terminating in a clavated tip.

Wings (when sitting) erect, and meeting upwards. (Flight diurnal.)

SPECIFIC CHARACTER.

Blackish Butterfly, with the upper wings spotted with yellow at the tips, the lower wings marked by a yellowish band, and beneath by a row of blue crescents.

This insect is a native of several parts of South America, and is subject to some slight variation as to colour: the plate represents it in its natural size.